

Trichoderma Secondary Metabolites Active on Plants and Fungal Pathogens

Francesco Vinale^{1,*}, Krishnapillai Sivasithamparam², Emilio L. Ghisalberti³, Sheridan L. Woo^{1,4}, Marco Nigro^{1,4}, Roberta Marra^{1,4}, Nadia Lombardi^{1,4}, Alberto Pascale^{1,4}, Michelina Ruocco¹, Stefania Lanzuise^{1,4}, Gelsomina Manganiello⁴ and Matteo Lorito^{1,4}

¹CNR – Istituto per la Protezione Sostenibile delle Piante (IPSP-CNR), Via Università 133, 80055 Portici, Italy

²School of Earth and Environment – Faculty of Natural and Agricultural Sciences, The University of Western Australia, W.A. 6009, Australia

³School of Chemistry and Biochemistry – Faculty of Life and Physical Sciences, The University of Western Australia, W.A. 6009, Australia

⁴Dipartimento di Agraria, Università degli Studi di Napoli ‘Federico II’, Via Università 100, 80055, Portici, Italy

Abstract: Beneficial microbes typically produce bioactive molecules that can affect the interactions of plants with their pathogens. Many secondary metabolites may also have antibiotic properties, which enable the producing microbe to inhibit and/or kill other microorganisms i.e. competing for a nutritional niche. Indeed, some of these compounds have been found to play an important role in the biocontrol of plant diseases by various beneficial microbes used world-wide for crop protection and bio-fertilization. In addition to direct toxic activity against plant pathogens, biocontrol-related metabolites may also increase disease resistance by triggering systemic plant defence activity, and/or enhance root and shoot growth. Fungi belonging to the *Trichoderma* genus are well known producers of secondary metabolites with a direct activity against phytopathogens and compounds that substantially affect the metabolism of the plant. The widescale application of selected metabolites to induce host resistance and/or to promote crop yield may become a reality in the near future and represents a powerful tool for the implementation of IPM strategies.

Keywords: Fungal interactions, plant protection, secondary metabolites, *Trichoderma*.

1. INTRODUCTION

Considerable crop losses incurred by plant diseases caused by phytopathogenic agents. Interest in biological control of fungal pathogens has increased recently in order to find alternatives to the use of chemicals. Synthetic pesticides are costly, pollute the environment and are potentially harmful to animals and humans. Furthermore, their repeated use promotes the development of chemically resistant pathogen strains.

The use of microbes for pest management in agriculture is one of the most effective strategies of biological control. The outcomes of using beneficial microbes are strain dependent and the advantages for the associated plant include: 1) establishment of an antagonistic microbial community in the rhizosphere; 2) suppression of pathogens; 3) overall improvement of plant health; 4) growth promotion; 5) increased nutrient availability and uptake, and 6) enhanced host resistance to both biotic and abiotic stresses [1-3].

The modes of action of beneficial microbes include: inhibition or parasitism of pathogens by using antibiotics often in combination with extracellular cell wall-degrading enzymes; competition for nutrients (i.e. iron, nitrogen or carbon) in colonization sites; stimulation of plant resistance mechanisms and development [4, 5].

Although not essential for their primary metabolic processes, microbes, and particularly fungi, produce various secondary metabolites (SMs), including compounds of industrial and economic relevance [6]. SMs are chemically different natural compounds of relatively low molecular weight (in most cases < 3 kDa), that are mainly produced by microorganisms and plants and typically associated to individual genera, species or strains. SMs are biosynthesized from primary metabolites in specialized pathways (i.e. polyketides or mevalonate pathways derived from Acetyl Coenzyme A, or amino acids) and some genes are clustered together. The expression of these genes appears to be induced by one or a few global regulators [6]. SMs show several biological activities possibly related to survival functions of the organism, such as competition against other micro- and macroorganisms, symbiosis, and metal transport.

In fungi, the production of SMs has been often correlated to specific stages of morphological differentiation, and

*Address correspondence to this author at the CNR – Istituto per la Protezione Sostenibile delle Piante (IPSP-CNR), Via Università 133, 80055 Portici, Italy; Tel: +39 081 2539338; Fax: +39 081 2539339; E-mail: f.vinale@ipp.cnr.it




Fig. (1). Chemical structure of: 1-6-pentyl- α -pyrone; 2. cytosporone S; 3,4,5,6,7. Koninginins A, B, D, E and G.

associated to the phase of active growth [7-9]. Their distribution ranges from a limited number of related species to single strains, indicating the absence of a universal function [7-9].

SMs exhibit several biological functions and play an important role in regulating interactions between organisms [10]. Some examples are phytotoxins (SMs produced by fungal pathogens that attack plants), mycotoxins (SMs produced by fungi that colonize crops capable of causing disease and death in humans and other animals), pigments (colored compounds also with antioxidant activity) and antibiotics (natural products capable of inhibiting or killing microbial competitors) [7-9]. However, biological activities are not necessarily confined to one specific group or single metabolites [10, 11].

Interestingly, some fungal secondary metabolites can modify the growth and the metabolism of plants, while others seem to target specific fungal processes such as sporulation and hyphal elongation [7]. Microbial metabolites are involved in several biological activities having important consequences in crop production. This review focuses on the main secondary metabolites produced by beneficial fungi and their roles in the complex interactions between the producing microbe and fungal pathogens or plants.

2. SECONDARY METABOLITES THAT ARE TOXIC TO OTHER FUNGI

Fungal SMs have been largely applied in human medicine, providing important drugs such as the antibiotics penicillin and cephalosporins, the immunosuppressant

cyclosporine and the antihypercholesterolemic agents lovastatin and compactin [12-17].

The involvement of toxic metabolites in plant disease and particularly in the interactions between beneficial and pathogenic fungi has been conclusively demonstrated by several studies [18]. Antibiotic production by antagonistic fungi is a well-documented phenomenon and often related to the strain biocontrol ability [19, 20].

Biocontrol isolates belonging to *Trichoderma* genus are well known producers of SMs that are toxic for phytopathogenic fungi [21, 22]. The following paragraphs report the most significant SMs isolated from *Trichoderma* spp. that have shown a significant antifungal activity.

Pyrones

The pyrone 6-pentyl-2H-pyran-2-one (6-pentyl- α -pyrone or 6PP – Fig. 1) is a metabolite commonly purified in the culture filtrate of different *Trichoderma* species (*T. viride*, *T. atroviride*, *T. harzianum*, *T. koningii*) and is responsible for the coconut aroma released by axenically developed colonies. 6PP has shown both *in vivo* and *in vitro* antifungal activities towards several plant pathogenic fungi and a strong relationship has been found between the biosynthesis of this metabolite and the biocontrol ability of the producing microbe [23, 24].

6PP biosynthesis in *T. atroviride* “is regulated by the G protein Tga1; other uncharacterized metabolites, however, are overproduced in tga1 mutants” [25]. The transcription factor Thctf1 also regulates the biosynthesis of 6PP in *T. harzianum*. Chromatographic and spectroscopic analyses




Fig. (2). Chemical structure of: 8 viridin; 9 viridiol; 10 harzianopyridone; 11 harzianic acid; 12 T22azaphilone.




Fig. (3). Chemical structure of: 13 Harzianolide; 14 T39butenolide; 15 deydro-harzianolide; 16 cerinolactone; 16 Dermadin; 18 trichoviridin; 19 homothallins.

showed that *Thctf1* null mutants did not produce two secondary metabolites derived from 6PP (6-[(1'R,2'S)-dihydroxypentyl]-2H-pyran-2-one and 6-[(1'S,2'R)-2'-propyloxiran-1-yl]-2H-pyran-2-one) and not previously described in the *Trichoderma* genus, that are present in wild-type culture filtrates [26, 27].

Another pyrone named cytosporone S (Figs. 1-2), recently isolated from a *Trichoderma* sp. strain, has been reported to have *in vitro* antibiotic activity against several bacteria and fungi [28].

Koninginins

Koninginins are complex pyranes isolated from *T. harzianum*, *T. koningii* and *T. aureoviride*. Koninginins A, B, D, E and G (Figs. 1; 3-7) showed *in vitro* antibiotic activity towards the take-all fungus *Gaeumannomyces graminis* var. *tritici* [29, 30]. "Koninginin D (22) also

inhibited the growth of other important soil-borne plant pathogens, such as *Rhizoctonia solani*, *Phytophthora cinnamomi*, *Pythium middletonii*, *Fusarium oxysporum* and *Bipolaris sorokiniana*" [31].

Viridins

The steroidal metabolite viridin (Figs. 2-8) is an antifungal compound isolated from diverse *Trichoderma* spp. (*T. koningii*, *T. viride*, *T. virens*) [32, 33]. This molecule prevents spore germination of *Botrytis allii*, *Colletotrichum lini*, *Fusarium caeruleum*, *Penicillium expansum*, *Aspergillus niger* and *Stachybotrys atra* [21, 34].

T. viride, *T. hamatum* and certain *Gliocladium* species produce viridiol (Figs. 2-9), a similar antifungal and phytotoxic metabolite for which the *in vivo* activity has been demonstrated [35, 36].




Fig. (4). Chemical structure of: 20 gliotoxin; 21 gliovirin; 22 chaetomin; 23 bigutol.




Fig. (5). Chemical structure of: 24 macrosphelide A; 25 3(2H)-benzofuranones A; 26 3(2H)-benzofuranones B; 27 chromanes A; 28 chromanes B; 29 chromanes C; 30 coniothyriomycin.

“A transcriptional comparison of wild type and a viridin and viridiol deficient *T. vires* mutant resulted in the identification of six genes similar to those involved in secondary metabolism of other fungi. Four of these genes (three cytochrome P450s and a cyclase) are located as a cluster that is associated with the production of viridin” [37].

Nitrogen Heterocyclic Compounds

Harzianopyridone (Figs. 2-10), a *T. harzianum* metabolite containing a penta-substituted pyridine ring system with a 2,3-dimethoxy-4-pyridinol pattern, is a potent antibiotic active against *Botrytis cinerea*, *R. solani* [38], *G. graminis* var. *tritici* and *Pythium ultimum* [39].

Recently, a new compound named harzianic acid and characterized by the presence of a pyrrolidindione ring system (Figs. 2-11) has been isolated from a *T. harzianum* strain. This tetramic acid derivative showed *in vitro* antibiotic activity against *Pythium irregulare*, *Sclerotinia sclerotiorum* and *R. solani* [40].

Azaphilones

The azaphilones are natural products containing a highly oxygenated bicyclic core and a chiral quaternary center. A new azaphilone, named T22azaphilone (Figs. 2-12), that showed *in vitro* a marked growth inhibition of several plant pathogens (*R. solani*, *P. ultimum* and *G. graminis* var. *tritici*) has been isolated from liquid culture of *T. harzianum* T22 [39].

Butenolides and Hydroxy-Lactones

Harzianolide (Figs. 3-13) and its derivatives, T39butenolide (Figs. 3-14) and dehydro-harzianolide (Figs. 3-15), have been isolated from different strains of *T. harzianum* [29, 39, 41, 42]. The *in vitro* antifungal activities of these SMs was demonstrated against several phytopathogenic agents [29, 39].

A novel hydroxy-lactone derivative, named cerinolactone (Figs. 3-16), has been isolated from culture filtrates of

T. cerinum. The isolated compound showed *in vitro* antifungal activity against *P. ultimum*, *R. solani* and *B. cinerea* [43].

Isocyano Metabolites

Trichoderma spp. also produce isocyano metabolites that have a characteristic 5-membered ring. However, the isolation and separation of these compounds is very difficult due to their instability [21]. Dermadin (Figs. 3-16') from *T. viride* [44, 45], *T. koningii* [46] and *T. hamatum* [47] is an antibiotic metabolite that was patented in 1971 [48]. The isonitrile trichoviridin (Figs. 3-18) isolated from *T. koningii* [46-49] and *T. viride* showed *in vitro* antibiotic properties [50]. Several dermadin and trichoviridin analogues have also been isolated. Interestingly, *T. koningii* produces various cyclopentenones isocyano metabolites named homothallins (Figs. 3-19) that affect the morphology of *Phytophthora* spp. [21].

Diketopiperazines

Gliotoxin (Figs. 4-20) and gliovirin (Figs. 4-21) are the two most important *Trichoderma* secondary metabolites belonging to this class of compounds. "P group strains of *Trichoderma* (*Gliocladium*) *virens* produce the antibiotic gliovirin which is active against *P. ultimum*, but not against *R. solani*. Strains of the Q group produce gliotoxin, which is very active against *R. solani*, but less against *P. ultimum*" [51]. In seedling bioassay tests, strains of the P group were able to effectively control *Pythium* damping off on cotton, while Q group strains gave better results towards the same disease caused by *R. solani* [52, 53]. These studies clearly indicate the potential role of antibiotic production in the biocontrol mechanism of the gliotoxin/gliovirin producers.

"The *T. virens* *veA* ortholog *vell* (VELVET protein Vell) is involved in regulation of gliotoxin biosynthesis, biocontrol activity and many other secondary metabolism-related genes" [54, 55]. Moreover, deletion of the 4-phosphopantetheinyl transferase in *T. virens* resulted in a mutant that failed to induce resistance in *Arabidopsis* and inhibit pathogenic fungi through the production of antibiotics [56].

Peptaibols

Peptaibols are linear peptides rich in non-proteinogenic amino acids (i.e. α -aminoisobutyric acid and isovaline), acetylated at the N-terminal group and the C-terminus is an amino alcohol (i.e. phenylalaninol, valinol, leucinol, isoleucinol or tryptophanol) [57]. Lorito *et al.* [58] "demonstrated that peptaibols inhibited β -glucan synthase activity in the host fungus, while acting synergistically with *T. harzianum* β -glucanases. The inhibition of glucan synthase prevented the reconstruction of the pathogen cell walls, thus facilitating the disruptive action of β -glucanases". The most widely known peptaibol is the *T. viride* alamethicin. The terms peptaibiome and peptaibiomics (peptide antibiotics or peptaibiotics) have been suggested to describe the analysis and study of all peptaibols expressed in an organism or tissue [57] using spectrometric methods, like LC/ESI-MSⁿ [59] or intact-cell MALDI-TOF [60]. Large multifunctional enzymes known as peptide synthetases assemble peptaibols (non-ribosomal biosynthesis) by the

multiple carrier thiotemplate mechanism from a remarkable range of precursors, which can be N-methylated, acylated or reduced [61].

The role of toxic metabolites during the fungus-fungus interaction has been demonstrated also for *Chaetomium globosum*. The ability of several *C. globosum* strains to produce chaetomin (Figs. 4-22) was correlated with the ability to suppress *Pythium* damping off of sugar beet in pasteurized soil. Activity of chaetomin on *P. ultimum* was 10 times higher than that of the *Trichoderma* diketopiperazine gliotoxin [62].

Two antifungal metabolites (particularly against *R. solani*) named bigutol (Figs. 4-23) and its derivative methylbigutol, were isolated from three isolates of the mycoparasite *Verticillium biguttatum* [63].

Coniothyrium minitans, a mycoparasite of *Sclerotinia sclerotiorum* and *S. cepivorum* sclerotia, produced four closely related metabolites able to inhibit fungal growth. The main compound, identified as macrosphelide A (Figs. 5-24), inhibited *in vitro* the mycelial growth of *S. sclerotiorum* and *S. cepivorum* also at very low concentrations [64].

Two 3(2H)-benzofuranones and three chromanes (Figs. 5; 25-29) were isolated from another strain of *C. minitans* when the fungus was grown in artificial medium where the only carbon source was sterilized and ground sclerotia of *S. sclerotiorum* [65].

Finally, the coniothyriomycin (Figs. 5-30) isolated from a *Coniothyrium* sp. showed remarkable fungicidal activities [66].

3. METABOLITES THAT INHIBIT BIOACTIVE MOLECULES MADE BY THE OTHER FUNGI

Regarding secondary metabolites produced by beneficial fungi that inhibit bioactive molecules made by other fungi, only few examples are reported in literature. Examples of "SMs inhibitors" eventually relevant for biocontrol applications include the following compounds that act towards phytopathogenic fungi at specific infection stages. They may be produced by certain antagonistic and endophytic fungi also in pure culture [67]. Flaviolin (Figs. 6-31) is a non-toxic inhibitor of conidium germination in *Magnaporthe grisea*, that was isolated from liquid cultures of an unidentified Ascomycete [67]. This substance accumulates as an oxidized intermediate product of melanin biosynthetic pathway, especially in the presence of inhibitors of the 1,3,6,8-tetrahydroxynaphthalene reductase such as the commercial fungicide tricyclazole. Similarly, another selective inhibitor of conidium germination in *M. grisea* that showed no effects on vegetative growth is tenuazonic acid (Figs. 6-32), obtained from cultures of an endophytic *Cladosporium* sp. [67].

Four glisoprenins were purified from cultures of the antagonistic fungi *Clonostachys rosea* as inhibitors of appressorium formation in *M. grisea*. The most active of them was glisoprenin C (Figs. 6-33) [68, 69].

Appressorium formation in *M. grisea* was also specifically inhibited by the neobulgarones A-F, six dimeric anthraquinone derivatives isolated from mycelium of the




Fig. (6). Chemical structure of: 31 flaviolin; 32 tenuazonic acid; 33 glisopenin C; 34 neobulgarone D; 35 scytalol D; 36 cerulenin.

ascomycete *Neobulgaria pura* with neobulgarone D, **34** in Fig. (6), the most active compound [70].

Several non-toxic natural products, like scytalol D (Fig. 6-35), from the anamorphic fungus *Scytalidium* sp. [71] and cerulenin (Fig. 6-36) from *Cephalosporium caeruleum* [72], inhibited melanin biosynthesis, a very important process for the formation of an effective fungal appressorium [67].

Detoxification of fungal toxins by beneficial microorganisms also represents an interesting possibility for disease management. Pure cultures of fungi able to detoxify mycotoxins have been obtained from complex microbial populations by using enrichment culture techniques [73]. In particular, *T. viride* and other fungal species were able to degrade aflatoxin B1 [74]. Moreover, *T. harzianum* hydrolases were able to degrade aflatoxin B1 (AFB1) and Ochratoxin A (OTA) *in vitro*. The mycotoxin content in corn flour inoculated with 100 ppb of AFB1 was reduced by up to 30% after treatment with fungal culture filtrates. The enzymatic mixture was separated by gel filtration and fractions with a molecular weight of approximately 100 kDa showed the maximum capacity of mycotoxin degradation [75].

In substrates with high C/N ratios, *T. virens* (formerly *Gliocladium virens*) produced a metabolite similar to the antibiotic viridin (Fig. 2-8), called viridiol (Fig. 2-9), that acts as a plant growth inhibitor [36]. This compound, isolated also from *T. hamatum*, inhibited the 5'-hydroxyaverantin dehydrogenase, an enzyme included in aflatoxin biosynthesis, thus reducing or completely blocking the production of this mycotoxin during the fungal interactions [76, 77].

Elad *et al.* [78] found that mycoparasitism or antibiosis were not the main biocontrol mechanisms of *T. harzianum* strain T39 against *B. cinerea*. These authors indicated that the antagonist interferes with the infection process by affecting the pathogen conidia in the early stages of the interaction. They suggest that T39 acts directly by inhibiting

B. cinerea hydrolytic enzymes, or indirectly by blocking plant responses that induce enzymatic activity in *B. cinerea*. Subsequently, Elad and Kapat [79] demonstrated that a *Trichoderma* protease is involved in the biocontrol of *B. cinerea* by degrading the hydrolytic enzymes required for infection by the pathogen.

4. METABOLITES THAT SUPPORT COMPETITION FOR NUTRIENTS

Competition for carbon, nitrogen and iron plays an important role during the interactions between beneficial and detrimental fungi, and is associated with the biocontrol mechanisms of non-pathogenic *Fusarium* and *Trichoderma* species [3].

"*Trichoderma* has a strong capacity to mobilize and take up soil nutrients, which makes it more efficient and competitive than many other soil microbes" [3]. This process could be related also to the production of organic acids, such as gluconic, citric and fumaric acids, which decrease soil pH and allow the solubilization of phosphates, micronutrients and mineral cations like iron, manganese and magnesium [3].

Iron is a mineral essential nutrient for numerous microorganisms, both bacteria and fungi [80]. "As a transition metal, redox properties of iron allow it to exist in two oxidation states, ferrous (Fe^{2+}) and ferric (Fe^{3+}) for the donation and acceptance of electrons, respectively" [80]. In the aerobic environment (with oxygen and neutral pH) iron exists mainly as Fe^{3+} and tends to form not soluble hydroxides and oxyhydroxides, making it not available for microbes growth [80].

Microorganisms that excrete siderophores (low-molecular weight; 200-1000 Da) are able to grow in natural environments poor in iron using residual not available iron [81, 82]. Most fungi produce various siderophores, which help the microbes to overcome adverse conditions.

Typically, extracellular and intracellular siderophores are found in fungi that transport or store ferric iron [81].

Secreted metabolites form extracellular Fe(III) complexes. “Then, the iron-charged siderophore is taken up by ferric-chelate-specific transporters or the siderophore-bound Fe(III) undergoes reduction to Fe(II), which is catalyzed by free extracellular or membrane-standing ferric-chelate reductases. If not already released extra cytoplasmatically, the iron has to be removed from the Fe-siderophore complex in the cytosol. This is mediated either by intracellular ferric-siderophore reductases or, in a few cases, by ferric-siderophore hydrolases” [83].

During the steps of iron utilization, the acquired metal is transferred through intracellular trafficking pathways, which may include diverse storage compartments in order to be directed to cofactor assembly systems and to final protein targeting [80].

Transport of siderophores is an energy-dependent process and is stereoselective, depending on recognition of the metal ion coordination geometry. In addition to iron transport, siderophores produced by microorganisms have other functions and effects, including enhancement of pathogenicity, storage of intracellular iron and suppression of microbial growth during the competition with other microbes [80].

The production of microbial siderophores can be beneficial to plants for two reasons: i) siderophores can solubilize iron unavailable for the plant; ii) siderophore production by non-pathogenic microorganisms can also suppress the growth of plant pathogens by depriving them of iron sources [84].

The majority of the fungal siderophores isolated so far belongs to hydroxamate class and can be divided into three structural families: fusarinines, coprogens and ferrichromes [85]. Fungi typically produce more than one siderophore, even if restricted to a particular family. However, a few cases have been reported of fungi able to synthesize siderophores from different structural families [86].

In order to study the importance of iron concentration for the activity of a *T. asperellum* strain (T34), Segarra *et al.* [87] analysed the effect of iron during the interaction of T34 with *Fusarium oxysporum* f.sp. *lycopersici* on tomato plants. In these experiments “Fe competition is one of the key factors for the biocontrol activity of T34 against the pathogen, as an increase in Fe concentration in the nutrient solution lead to the suppression of T34 siderophore synthesis and thus to the inhibition of Fe competition” [87]. Moreover, T34-treated plants significantly enhanced plant growth and development as compared to control plants at high level of Fe (1,000 μ M), even though T34 did not reduce the Fe content in leaves or stems. In another work T34 increased the Fe concentration in the aerial parts of lupin plants grown on a calcareous medium. These results clearly indicate a role of siderophores during the interaction with the pathogen and the plants, although no siderophores have been isolated and characterized so far from the culture filtrate of this beneficial microbe [88].

Anke *et al.* [86] isolated siderophores from all different structural families simultaneously from a culture filtrate of

Trichoderma spp. In particular, the culture filtrate of this fungus obtained in iron deficiency condition contained coprogen, coprogen B, fusarinine C and ferricrocin (Figs. 7; 37-40).

In a recent work a new method for the analysis of iron-chelating metabolites using LC-HRMS was used to detect extracellular siderophores produced by 10 different *Trichoderma* strains. *T. harzianum* produced the highest number of siderophores and did not have any unique compounds; while, *T. reesei* biosynthesized one cis-fusarinine as the major siderophore and three others that were present only in *T. harzianum*. The data suggest “that the high diversity of siderophores produced by *Trichoderma* spp. might be the result of further modifications of the non-ribosomal peptide synthetase (NRPS) products and not due to diverse NRPS-encoding genes” [89].

Recently, the ability of the *Trichoderma* SM harzianic acid (Figs. 2-11) to bind with a good affinity essential metals such as Fe³⁺ has been demonstrated [90].

5. METABOLITES INVOLVED IN BENEFICIAL FUNGUS-PLANT INTERACTIONS

SMs that Increase Systemic Resistance

Several metabolites produced by beneficial fungi are involved in the induction of plant resistance, such as: i) “proteins with enzymatic activity, i.e. xylanase” [91]; ii) “avirulence-like gene products able to induce defence reactions in plants” [2]; and iii) “low-molecular-weight compounds released from either fungal or plant cell walls” by specific enzyme activities [2, 92, 93]. “Some of the low-molecular-weight degradation products released from fungal cell walls have been purified and characterized, and found to consist of short oligosaccharides comprised of two types of monomers, with and without an amino acid residue” [92, 93]. “These compounds elicited a reaction in the plant when applied to leaves or when injected into root or leaf tissues. Further, they also stimulated the biocontrol ability of fungi such as *Trichoderma* by activating the mycoparasitic gene expression cascade” [92].

Some *Trichoderma* SMs may act as elicitors of plant defence mechanisms against pathogens. A reduction of disease symptoms on tomato and canola seedlings treated in particular with 6PP (Fig. 1) and inoculated, respectively, with the pathogens *B. cinerea* or *Leptosphaeria maculans* has been reported [94]. Moreover, “soil drench applications of 6PP four days before inoculation with *Fusarium moniliforme* showed considerable suppression of seedling blight and substantial plant growth promotion, compared with the untreated control” [95]. Application of 6PP on maize seedlings distinctly enhanced the activities of peroxidase, polyphenoloxidase and β -1,3-glucanase in both shoot and root tissues indicating an induction of defence responses in maize plants [95].

Peptaibols are another class of plant defence elicitors produced by *Trichoderma*. Application of alamethicin, a long sequence peptaibol with a 20-residue produced by *T. viride*, induced defence responses in *Phaseolus lunatus* (lima bean) [96] and *Arabidopsis thaliana* [97]. Moreover,




Fig. (7). Chemical structure of: 37 coprogen; 38 coprogen B; 39 fusarinine C; 40 ferricrocin.

disruption of the non-ribosomal peptide synthetases (NRPS) gene, *tex1*, resulted in the loss of production of all forms of 18-residue peptaibols in *T. virens* which corresponded to a significant reduction of the ability of the fungus to induce systemic resistance responses [98].

Recently Mukherjee *et al.* [99] demonstrated that mutation in one of the polyketide synthase/non-ribosomal peptide synthetases “(PKS/NRPS) hybrid genes reduces the ability of *T. virens* to induce the defence response gene *pal* (phenylalanine ammonia lyase), suggesting a putative role for the associated metabolite” [99] (derived from polyketide pathway) product in induced systemic resistance. These results provide evidence that a PKS/NRPS hybrid enzyme responsible for the metabolite production is involved in

Trichoderma–plant interactions resulting in induction of defence response in maize.

SMs that Enhance Growth, Development and Yield

Many beneficial fungi including some *Trichoderma* species, can increase plant growth, development and yield also in the absence of pathogens [100-104].

Several strains belonging to the *Trichoderma* genus, including some used world-wide in agriculture, have been found to be able to stimulate plant development, especially at the root level (i.e. formation of more lateral roots) by activating an auxin-dependent mechanism [105] and/or producing indole-3-acetic acid (IAA) or auxin analogues [106].




Fig. (8). Chemical structure of: 41, 42, 43, 44 trichocaranes A – D; 45 cyclonerodiol; 46 trichosetin; 47 5-hydroxy-1-(3-methyl-3-buten-1-ynyl)-7-oxabicyclo[4.1.0]-hept-3-en-2-one.

Positive effect on plant development have been demonstrated for several *Trichoderma* SMs [107]. Koninginins (Fig. 1; 3-7), 6PP (Fig. 1), trichocaranes A – D (Fig. 8; 41-44), harzianopyridone (Fig. 2-9), cyclonerodiol (Fig. 8-45), harzianolide (Fig. 3-13) and harzianic acid (Fig. 2-11) are examples of isolated compounds that affect plant growth in a concentration dependent manner [30, 108-114].

A dual culture fungus/plant provides a simple method of establishing their interaction and allows the isolation of metabolites induced by one of the two components. Interestingly, dual culture of *T. harzianum* and calli of *Catharathus roseus* produced an antimicrobial metabolite called trichosetin (Fig. 8-46), that was absent in single cultures. This compound was probably produced by the fungus [115]. Trichosetin affected negatively growth and development of numerous plant species [116] and is a *N*-desmethyl analogue of equisetin, a tetramic acid isolated from *Fusarium equiseti* [117].

A novel metabolite, named cerinolactone (Fig. 3-16), has been isolated and characterized *T. cerinum* and was able to positively alter the growth of tomato seedlings three days after treatment [43].

A sterile dark ectotrophic fungus isolated from roots of *Neurachne alopecuroidea* produced 5-hydroxy-1-(3-methyl-3-buten-1-ynyl)-7-oxabicyclo[4.1.0]-hept-3-en-2-one in liquid cultures (Fig. 8-47). The metabolite showed antimicrobial activity against some plant pathogens and improve plant growth and development, similarly to what found for the ectotrophic fungus applied *in vivo* [118].

The dose-effect response of plant growth to secondary metabolites produced by beneficial fungi clearly deserves further investigation. These metabolites may possibly act as auxin-like molecules, which have a positive effect at low concentrations while having an inhibitory effect at higher doses. For instance, an hormone activity was detected on

etiolated pea stems treated with harzianolide and 6PP. These compounds also affected the growth of tomato and canola seedlings in a manner depending on the concentration and/or the application method used [94].

CONCLUSION

Hundreds of SMs produced by beneficial fungi have been isolated and characterized. Here we focused only on the compounds produced by fungal microorganisms that have been involved in the interactions with phytopathogenic agents and/or plants having a positive outcome for agriculture.

In terms of a sheer number, fungal SMs with a direct antibiotic activity against plant pathogens have been mainly isolated from biocontrol strains of the genus *Trichoderma*. Even though many SMs are known, elite strains usually produce only a few main SMs. The quality and the quantity of SMs synthesized depend on: i) the compound considered; ii) the species and the strain; iii) the occurrence of other microorganisms; iv) the equilibrium among elicited biosynthesis and biotransformation rate; v) the growth conditions. Further, in some cases, the biocontrol agent was able to modulate the production of toxic SMs according to the presence or the absence of the target pathogen [119].

Interestingly, SMs by beneficial fungi may also be involved in biocontrol mechanisms through the inhibition of bioactive products produced by fungal pathogens. Examples of metabolites that inhibit other fungal products or bioactive molecules include compounds able to detoxify or inhibit the biosynthesis of mycotoxins, which is very harmful for humans and animals. Further studies on this interesting topic may produce some new biotechnological tools to effectively reduce contamination and losses of human food and animal feed.

It is well recognized that biocontrol fungi, such as selected agents of *Trichoderma* spp., are able to produce

compounds with multiple activities, including direct/indirect toxic effects against plant pathogens, plant defence induction or growth promotion [3]. Several fungal species, as well as numerous chemically different substances, have been found to be able to modify plant development and crop yield. An hormone-like effect has been proposed for some *Trichoderma* SMs and specific antimicrobial compounds having this characteristics have been detected in plant-fungus cultures. In fact, treatment with *Trichoderma* metabolites produces extensive changes of the plant expressome, proteome and metabolome, by acting on specific pathways involved in the synthesis of major hormones, resistance to biotic/abiotic stresses and nutrient uptake [107]. These recent findings have suggested new strategies for the development of novel bioformulates based on microbial metabolites alone or in combination with live microbes, in order to maximize the beneficial effects and reduce the risks associated with the release of microorganisms into the environment.

The current techniques, such as metabolomics and expressomics, could provide novel information about the molecular factors involved in the complex interactions occurring between plants, beneficial microbes and phytopathogenic agents. Nevertheless, it is important to continue with the identification of natural compounds that are not produced under usual laboratory conditions [120]. Therefore, novel techniques developed recently need to be used in order to select *Trichoderma* strains for biocontrol that are able to produce only or mainly “beneficial” metabolites (i.e. antibiotics not toxic to plants and their consumers and/or acting as plant growth promoters and/or inducers of disease resistance) [120].

The application of selected metabolites to induce host resistance and/or to promote crop yield may be an interesting alternative to chemicals. Regardless, further studies aimed at conclusively determining the nature and the fate of mixtures of SMs released in the soil or the phyllosphere by beneficial fungi applied as biocontrol or fertilization agents by employing “inundative methods”, are still needed.

CONFLICT OF INTEREST

The authors confirm that this article content has no conflict of interest.

ACKNOWLEDGEMENTS

Work by the authors FV and MR was supported by a dedicated grant from the Italian Ministry of Economy and Finance to the National Research Council for the project "Innovazione e Sviluppo del Mezzogiorno - Conoscenze Integrate per Sostenibilità ed Innovazione del Made in Italy Agroalimentare - Legge n. 191/2009". Work by the authors FV, MR, SW, ML, MN, RM, NL, AP, SL and GM has been supported by the following projects: Calabria Region APQ Projects “S.Re.Va.Pr.O.”, Mis. 124, Projects “CA.SVI.PR.OLI.” and “MI.P.RE.VEGE.”; PRIN MIUR 2008 - prot. 2008SNPNC2 - prot. 2008WKPAWW; FIRB 2002 prot. RBNE01K2E7; PRIN 2003 prot. 2003070719-003; MIUR project SICURA; MIUR PON project no. DD12935 of 02/08/2002; MIUR PON project no. DD1219 of 05/10/2004; MIUR PON project no. DD1801 of 31/12/2004;

MIUR PON project no. DD29156 of 20/04/2011; EU TRICHOEST QLK3-2002-02032; EU 2E-BCA2.

REFERENCES

- [1] Harman GE. Myths and dogmas of biocontrol: changes in perceptions derived from research on *Trichoderma harzianum* T-22. *Plant Dis* 2000; 84: 377-93.
- [2] Harman GE, Howell CR, Viterbo A, Chet I, Lorito M. *Trichoderma* species - opportunistic, avirulent plant symbionts. *Nat Rev Microbiol* 2004; 2: 43-56.
- [3] Vinale F, Sivasithamparam K, Ghisalberti EL, Marra R, Woo SL, Lorito M. *Trichoderma*-plant-pathogen interactions. *Soil Biol Biochem* 2008; 40: 1-10.
- [4] Howell CR. Mechanisms employed by *Trichoderma* species in the biological control of plant diseases: the history and evolution of current concepts. *Plant Dis* 2003; 87: 4-10.
- [5] Whipps JM. Microbial interactions and biocontrol in the rhizosphere. *J Exp Bot* 2001; 52: 487-511.
- [6] Herbert RB. The biosynthesis of secondary metabolites. 2nd Ed. London: Chapman & Hall; Horsfall 1989.
- [7] Keller NP, Turner G, Bennett JW. Fungal secondary metabolism — from biochemistry to genomics. *Nat Rev Microbiol* 2005; 3: 937-47.
- [8] Demain AL, Fang A. The natural functions of secondary metabolites. *Adv Biochem Eng Biot* 2000; 69: 1-39.
- [9] Chiang Y, Lee K, Sanchez JF, Keller NP, Wang CCC. Unlocking fungal cryptic natural products. *Nat Prod Commun* 2009; 4: 1505-10.
- [10] Hanson JR. Natural products: the secondary metabolites. Vol. 17 Ed. Cambridge: Royal Society of Chemistry 2003.
- [11] Hanson JR. The chemistry of fungi. Ed. Cambridge: Royal Society of Chemistry 2008.
- [12] Fleming, A. On the antibacterial action of cultures of a *Penicillium*, with special reference to their use in the isolation of *B. influenzae*. *Brit J Exp Pathol* 1929; 10: 226-36.
- [13] Ruegger A, Kuhn M, Lichti H. Cyclosporin A, a peptide metabolite from *Trichoderma polysporum* (Link ex Pers.) Rifai, with a remarkable immunosuppressive activity. *Helv Chim Acta* 1976; 59: 1075-92.
- [14] Endo A, Kuroda M, Tsujita Y. ML-236A, ML-236B, and ML-236C, new inhibitors of cholesterologenesis produced by *Penicillium citrinium*. *J Antibiot* 1976; 29: 1346-8.
- [15] Endo A, Monacolin K. A new hypocholesterolemic agent produced by a *Monascus* species. *J Antibiot* 1979; 32: 852-4.
- [16] Alberts AW. Mevinolin: a highly potent competitive inhibitor of hydroxymethylglutaryl-coenzyme A reductase and a cholesterol-lowering agent. *Proc Natl Acad Sci USA* 1980; 77: 3957-61.
- [17] Sammartino G, Trosino O, di Lauro AE, Amato M, Cioffi A. Use of piezosurgery in management of surgical dental implant complication: a case report. *Implant Dent* 2011; 20: e1-6.
- [18] Fravel DR. Role of antibiosis in the biocontrol of plant diseases. *Annu Rev Phytopathol* 1988; 26: 75-91.
- [19] Howell CR. The role of antibiosis in biocontrol. In: *Trichoderma and Gliocladium*, Vol. 2, Harman GE, Kubicek CP (Eds). Taylor and Francis Ltd: London UK 1998; pp. 139-91.
- [20] Sivasithamparam K, Ghisalberti EL. Secondary metabolism in *Trichoderma* and *Gliocladium*. In: *Trichoderma and Gliocladium*, Vol. 1, Harman GE, Kubicek CP (Eds). Taylor and Francis Ltd: London UK 1998; pp. 139-91.
- [21] Reino JL, Guerrero RF, Hernández-Galán R, Collado IG. Secondary metabolites from species of the biocontrol agent *Trichoderma*. *Phytochem Rev* 2008; 7: 89-123.
- [22] Ghisalberti EL. Anti-infective agents produced by the hyphomycetes genera *Trichoderma* and *Gliocladium*. *Curr Med Chem* 2002; 1: 343-74.
- [23] Scarselletti R, Faull JL. *In Vitro* activity of 6-pentyl-a-pyrone, a metabolite of *Trichoderma harzianum*, in the inhibition of *Rhizoctonia solani* and *Fusarium oxysporum* f. sp. *lycopersici*. *Mycol Res* 1994; 98:1207-09.
- [24] Worasatit N, Sivasithamparam K, Ghisalberti EL, Rowland C. Variation in pyrone production, pectic enzymes and control of rhizoctonia root rot of wheat among single-spore isolates of *Trichoderma koningii*. *Mycol Res* 1994; 98: 1357-63.

- [25] Reithner B, Brunner K, Schuhmacher R, *et al.* The G protein alpha subunit Tgal of *Trichoderma atroviride* is involved in chitinase formation and differential production of antifungal metabolites. *Fungal Genet Biol* 2005; 42: 749-60.
- [26] Rubio MB, Hermosa R, Reino JL, Collado IG, Monte E. Thctf1 transcription factor of *Trichoderma harzianum* is involved in 6-pentyl-2H-pyran-2-one production and antifungal activity. *Fungal Genet Biol* 2009; 46: 17-27
- [27] Daoubi M, Pinedo-Rivilla C, Rubio MB, *et al.* Hemisynthesis and absolute configuration of novel 6-pentyl-2H-pyran-2-one derivatives from *Trichoderma* spp. *Tetrahedron* 2009; 69: 4834-40.
- [28] Takahiro I, Nonaka K, Suga T, Masuma R, Omura S, Shiomi K. Cytosporone S with antimicrobial activity, isolated from the fungus *Trichoderma* sp. FKI-6626. *Bioorg Med Chem Lett* 2013; 23: 679-81.
- [29] Almassi F, Ghisalberti EL, Narbey MJ, Sivasithamparam K. New antibiotics from strains of *Trichoderma harzianum*. *J Nat Prod* 1991; 54: 396-402.
- [30] Ghisalberti EL, Rowland CY. Antifungal metabolites from *Trichoderma harzianum*. *J Nat Prod* 1993; 56: 1799-804.
- [31] Dunlop RW, Simon A, Sivasithamparam K, Ghisalberti EL. An antibiotic from *Trichoderma koningii* active against soilborne plant pathogens. *J Nat Prod* 1989; 52: 67-74.
- [32] Golder WS, Watson TR. Lanosterol derivatives as precursors in the biosynthesis of viridin. Part 1. *J Chem Soc Perkin Trans* 1980; 1: 422-5.
- [33] Singh S, Dureja P, Tanwar RS, Singh A. Production and antifungal activity of secondary metabolites of *Trichoderma virens*. *Pestic Res J* 2005; 17: 26-9.
- [34] Brian PW, McGowan JC. Viridin: a highly fungistatic substance produced by *Trichoderma viride*. *Nature* 1945; 156: 144-5.
- [35] Moffatt JS, Bu'Lock JD, Yuen TH. Viridiol, a steroid-like product from *Trichoderma viride*. *J Chem Soc Chem Commun* 1969; 14: 839.
- [36] Howell CR, Stipanovic RD. Effect of sterol biosynthesis inhibitors on phytotoxin (viridiol) production by *Gliocladium virens* in culture. *Phytopathology* 1994; 84: 969-72.
- [37] Mukherjee M, Horwitz BA, Sherkhane PD, Hadar R, Mukherjee PK. A secondary metabolite biosynthesis cluster in *Trichoderma virens*: evidence from analysis of genes underexpressed in a mutant defective in morphogenesis and antibiotic production. *Curr Genet* 2006; 50: 193-202.
- [38] Dickinson JM, Hanson JR, Hitchcock PB, Claydon N. Structure and biosynthesis of harzianopyridone, an antifungal metabolite of *Trichoderma harzianum*. *J Chem Soc Perkin Trans* 1989; 1: 1885-7.
- [39] Vinale F, Marra R, Scala F, Ghisalberti EL, Lorito M, Sivasithamparam K. Major secondary metabolites produced by two commercial *Trichoderma* strains active against different phytopathogens. *Lett App Microbiol* 2006; 43: 143-8.
- [40] Vinale F, Flematti G, Sivasithamparam K, *et al.* Harzianic acid, an antifungal and plant growth promoting metabolite from *Trichoderma harzianum*. *J Nat Prod* 2009; 72: 2032-5.
- [41] Claydon N, Hanson JR, Truneh A, Avent AG. Harzianolide, a butenolide metabolite from cultures of *Trichoderma harzianum*. *Phytochemistry* 1991; 30: 3802-3.
- [42] Ordentlich A, Wiesman Z, Gottlieb HE, Cojocar M, Chet I. Inhibitory furanone produced by the biocontrol agent *Trichoderma harzianum*. *Phytochemistry* 1992; 31: 485-6.
- [43] Vinale F, Arjona GI, Nigro M, *et al.* Cerinolactone, a hydroxy-lactone derivative from *Trichoderma cerinum*. *J Nat Prod* 2012; 75: 103-6.
- [44] Pyke TR, Dietz A. U-21,963, a new antibiotic. I. Discovery and biological activity. *Appl Microbiol* 1966; 14: 506-10
- [45] Meyer CE. U-21,963, a new antibiotic. II. Isolation and characterization. *Appl Microbiol* 1966; 14: 511-2.
- [46] Tamura A, Kotani H, Naruto S. Trichoviridin and dermadin from *Trichoderma* sp. TK-1. *J Antibiot* 1975; 28: 161-2.
- [47] Fujiwara A, Okuda T, Masuda S, *et al.* Isonitrile antibiotics, a new class of antibiotics with an isonitrile group. I. Fermentation, isolation and characterization of isonitrile antibiotics. *Agric Biol Chem* 1982; 46: 1803-9.
- [48] Coats JH, Meyer CE, Pyke TR, inventor; Antibiotic dermadin. US Patent 3627882, 14 Dec 1971.
- [49] Nobuhara M, Tazima H, Shudo K, Itai A, Okamoto T, Itaka Y. A fungal metabolite, novel isocyanate epoxide. *Chem Pharm Bull* 1976; 24: 832-4.
- [50] Yamano T, Hemmi S, Yamamoto I, Tsubaki K, inventor; Trichoviridin, a new antibiotic. JP Patent 45015435, 29 May 1970.
- [51] Howell CR. Selective isolation from soil and separation *in vitro* of P and Q strains of *Trichoderma virens* with differential media. *Mycologia* 1999; 91: 930-4.
- [52] Howell CR. Biological control of *Pythium* damping off of cotton with seed coating preparations of *Gliocladium virens*. *Phytopathology* 1991; 81: 738-41.
- [53] Howell CR, Stipanovic RD, Lumsden RD. Antibiotic production by strains of *Gliocladium virens* and its relation to the biocontrol of cotton seedling diseases. *Biocontrol Sci Techn* 1993; 3: 435-41.
- [54] Mukherjee PK, Kenerley CM. Regulation of morphogenesis and biocontrol properties in *Trichoderma virens* by a VELVET protein, Vel1. *Appl Environ Microbiol* 2010; 76: 2345-52.
- [55] Mukherjee PK, Horwitz BA, Herrera-Estrella A, Schmol M, Kenerley CM. *Trichoderma* research in the genome era. *Annu Rev Phytopathol* 2013; 51:105-29.
- [56] Velazquez-Robledo R, Contreras-Cornejo H, Macias-Rodriguez LI, *et al.* Role of the 4-phosphopantetheinyl transferase of *Trichoderma virens* in secondary metabolism, and induction of plant defense responses. *Mol Plant-Microbe Interact* 2011; 24: 1459-71.
- [57] Daniel JF, Filho ER. Peptaibols of *Trichoderma*. *Nat Prod Rep* 2007; 24: 1128-41.
- [58] Lorito M, Farkas V, Rebuffat S, Bodo B, Kubicek CP. Cell wall synthesis is a major target of mycoparasitic antagonism by *Trichoderma harzianum*. *J Bacteriol* 1996; 178: 6382-5.
- [59] Degenkolb T, Gräfenhan T, Berg A, Nirenberg HI, Gams W, Brückner H. Peptaibiotics: screening for polypeptide antibiotics (Peptaibiotics) from plant-protective *Trichoderma* species. *Chem Biodivers* 2006; 3: 593-610.
- [60] Neuhof T, Dieckmann R, Druzhinina IS, Kubicek CP, von Dohren H. Intact-cell MALDI-TOF mass spectrometry analysis of peptaibol formation by the genus *Trichoderma/Hypocrea*: can molecular phylogeny of species predict peptaibol structures? *Microbiology* 2007; 153: 3417-37.
- [61] Szekeres A, Leitgeb B, Kredics L, *et al.* Peptaibols and related peptaibiotics of *Trichoderma*. *Acta Microbiol Immunol Hung* 2005; 52: 137-68.
- [62] Di Pietro A, Gut-Rella M, Pachlatko JP, Schwinn FJ. Role of antibiotics produced by *Chaetomium globosum* in biocontrol of *Pythium ultimum*, a causal agent of damping-off. *Phytopathology* 1992; 82: 131-5.
- [63] Morris RAC, Ewing D, Whipps JM, Coley-Smith JR. Antifungal hydroxymethyl phenols from the mycoparasite *Verticillium biguttatum*. *Phytochemistry* 1995; 39: 1043-8.
- [64] McQuilken MP, Gemmill J, Hill RA, Whipps JM. Production of macrophelide A by the mycoparasite *Coniothyrium minitans*. *FEMS Microbiol Lett* 2003; 219: 27-31.
- [65] Machida K, Trifonov LS, Ayer WA, *et al.* 3(2H)-Benzofuranones and chromanes from liquid cultures of the mycoparasitic fungus *Coniothyrium minitans*. *Phytochemistry* 2001; 58: 173-7.
- [66] Krohn K, Franke C, Jones PG, Aust HJ, Draeger S, Schulz B. Biologically active metabolites of fungi. I. Isolation, synthesis, and biological activity of coniothyriomycin as well as bioassay of analogous open-chain imides. *Liebigs Ann Chem* 1992; 8: 789-98.
- [67] Thines E, Anke H, Weber RW. Fungal secondary metabolites as inhibitors of infection-related morphogenesis in phytopathogenic fungi. *Mycol Res* 2004; 108:14-25.
- [68] Thines E, Eilbert F, Sterner O, Anke H. Glisoprenin A, an inhibitor of the signal transduction pathway leading to appressorium formation in germinating conidia of *Magnaporthe grisea* on hydrophobic surfaces. *FEMS Microbiol Lett* 1997; 151: 219-24.
- [69] Sterner O, Thines E, Eilbert F, Anke H. Glisoprenins C, D and E, new inhibitors of appressorium formation in *Magnaporthe grisea* from cultures of *Gliocladium roseum*. *J Antibiot* 1998; 51: 228-31.
- [70] Eilbert F, Anke H, Sterner O. Neobulgarones A-F from cultures of *Neobulgaria pura*, new inhibitors of appressorium formation in *Magnaporthe grisea*. *J Antibiot* 2000; 53: 1123-9.
- [71] Thines E, Anke H, Sterner O. Scytalols A, B, C, and D and other modulators of melanin biosynthesis from *Scytalidium* sp. 36-93. *J Antibiot* 1997; 51: 387-93.

- [72] Sano Y, Nomura S, Kamio Y, Omura S, Hata T. Studies on cerulenin. 3. Isolation and physicochemical properties of cerulenin. *J Antibiot* 1967; 20: 344-8.
- [73] Karlovsky P. Biological detoxification of fungal toxins and its use in plant breeding, feed and food production. *Nat Toxins* 1999; 7: 1-23.
- [74] Mann R, Rehm HJ. Degradation products from aflatoxin B1 by *Corynebacterium rubrum*, *Aspergillus niger*, *Trichoderma viride*, and *Mucor ambiguus*. *Eur J Appl Microbiol* 1976; 2: 297-306.
- [75] Aloj V, Vinale F, Woo S, *et al.* Use of a *Trichoderma* spp. enzyme mixture to increase feed digestibility and degrade mycotoxins. *J Plant Pathol* 2009; 91: S4.45-96.
- [76] Sakuno E, Yabe K, Hamasaki T, Nakajima HA. New inhibitor of 5'-hydroxyaverantin dehydrogenase, an enzyme involved in aflatoxin biosynthesis, from *Trichoderma hamatum*. *J Nat Prod* 2000; 63: 1677-8.
- [77] Wipf P, Kerekes AD. Structure reassignment of the fungal metabolite TAEMC161 as the phytotoxin viridiol. *J Nat Prod* 2003; 66: 716-871.
- [78] Elad Y. Mechanisms involved in the biological control of *Botrytis cinerea* incited diseases. *Eur J Plant Pathol* 1996; 102: 719-32.
- [79] Elad Y, Kapat A. 1999. The role of *Trichoderma harzianum* protease in the biocontrol of *Botrytis cinerea*. *Eur J Plant Pathol* 1999; 105:177-89.
- [80] Miethke M. Molecular strategies of microbial iron assimilation: from high-affinity complexes to cofactor assembly systems. *Metallomics* 2013; 5: 15-28.
- [81] Winkelmann G. Ecology of siderophores with special reference to the fungi. *Biometals* 2007; 20: 379-92.
- [82] Johnson L. Iron and siderophores in fungal-host interactions. *Mycol Res* 2008; 112: 170-83.
- [83] Miethke M, Marahiel MA. Siderophore-based iron acquisition and pathogen control. *Microbiol Mol Biol R* 2007; 71: 413-51.
- [84] Leong J. Siderophores: their biochemistry and possible role in the biocontrol of plant pathogens. *Annu Rev Phytopathol* 1986; 24: 187-209.
- [85] Renshaw JC, Robson GD, Trinci API, *et al.* Fungal siderophores: structures, functions and applications. *Mycol Res* 2002; 106: 1123-42.
- [86] Anke H, Kinn J, Bergquist KE, Sterner O. Production of siderophores by strains of the genus *Trichoderma* - isolation and characterization of the new lipophilic coprogen derivative, palmitoylcoprogen. *Biometals* 1991; 4: 176-80.
- [87] Segarra G, Casanova E, Avilés M, Trillas I. *Trichoderma asperellum* strain T34 controls Fusarium wilt disease in tomato plants in soilless culture through competition for iron. *Microb Ecol* 2010; 59: 141-9.
- [88] de Santiago A, Quintero JM, Avilés M, Delgado A. Effect of *Trichoderma asperellum* strain T34 on iron nutrition in white lupin. *Soil Biol Biochem* 2009; 41: 2453-9.
- [89] Lehner SM, Atanasova L, Neumann NK, *et al.* Isotope-assisted screening for iron-containing metabolites reveals high diversity among known and unknown siderophores produced by *Trichoderma* spp. *Appl Environ Microbiol* 2013; 79: 18-31.
- [90] Vinale F, Nigro N, Sivasithamparam K, *et al.* Harzianic acid: a novel siderophore from *Trichoderma harzianum*. *FEMS Microbiol Lett* 2013; (In Press doi: 10.1111/1574-6968.12231).
- [91] Lotan T, Fluhr R. Xylanase, a novel elicitor of pathogenesis-related proteins in tobacco, uses a nonethylene pathway for induction. *Plant Physiol* 1990; 93: 811-7.
- [92] Woo SL, Scala F, Ruocco M, Lorito M. The molecular biology of the interactions between *Trichoderma* spp., phytopathogenic fungi, and plants. *Phytopathology* 2006; 96: 181-5.
- [93] Woo SL, Lorito M. In: Vurro M, Gressel J, Ed. Novel biotechnologies for biocontrol agent enhancement and management. Amsterdam: IOS, Springer Press. 2007; 107-30.
- [94] Vinale F, Sivasithamparam K, Ghisalberti EL, *et al.* A novel role for *Trichoderma* secondary metabolites in the interactions with plants. *Physiol Mol Plant P* 2008; 72: 80-6.
- [95] El-Hasan A, Buchenauer H. Actions of 6-pentyl-alpha-pyrone in controlling seedling blight incited by *Fusarium moniliforme* and inducing defence responses in maize. *J Phytopathol* 2009; 157: 697-707.
- [96] Engelberth J, Koch T, Schuler G, Bachmann N, Rechtenbach J, Boland W. Ion channel-forming alamethicin is a potent elicitor of volatile biosynthesis and tendrils coiling. Cross talk between jasmonate and salicylate signaling in lima bean. *Plant Physiol* 2000; 125: 369-77.
- [97] Chen F, D'Auria JC, Tholl D, *et al.* An *Arabidopsis thaliana* gene for methylsalicylate biosynthesis, identified by a biochemical genomics approach, has a role in defence. *Plant J* 2003; 36: 577-88.
- [98] Viterbo A, Wiest A, Brotman Y, Chet I, Kenerley C. The 18mer peptaibols from *Trichoderma virens* elicit plant defence responses. *Mol Plant Pathol* 2007; 8: 737-46.
- [99] Mukherjee PK, Buensanteai N, Moran-Diez ME, Druzhinina IS, Kenerley CM. Functional analysis of non-ribosomal peptide synthetases (NRPSs) in *Trichoderma virens* reveals a polyketide synthase (PKS)/NRPS hybrid enzyme involved in the induced systemic resistance response in maize. *Microbiology* 2012; 158: 155-65.
- [100] Chang YC, Baker R, Kleifeld O, Chet I. Increased growth of plants in the presence of the biological control agent *Trichoderma harzianum*. *Plant Dis* 1986; 70: 145-8.
- [101] Harris AR. Plant growth promotion by binucleate *Rhizoctonia* and bacterial isolates in monoxenic cultures. *Microbiol Res* 1999; 154: 71-4.
- [102] Shivanna MB, Meera MS, Hyakumachi M. Role of root colonization ability of plant growth promoting fungi in the suppression of take all and common root rot of wheat. *Crop Protec* 1996; 15: 497-504.
- [103] Windham MT, Elad Y, Baker R. A mechanism for increased plant growth induced by *Trichoderma* spp. *Phytopathology* 1986; 76: 518-21.
- [104] Wulff EG, Pham ATH, Chérif M, Rey P, Tirilly Y, Hockenull J. Inoculation of cucumber roots with zoospores of mycoparasitic and plant pathogenic *Pythium* species: differential zoospore accumulation, colonization ability and plant growth response. *Europ J Plant Pathol* 1998; 104: 69-76.
- [105] Contreras-Cornejo HA, Macías-Rodríguez L, Cortés-Penagos C, López-Bucio J. *Trichoderma virens*, a plant beneficial fungus, enhances biomass production and promotes lateral root growth through an auxin-dependent mechanism in *Arabidopsis*. *Plant Physiol* 2009; 149: 1579-92.
- [106] Hoyos-Carvajal L, Orduz S, Bissett J. Growth stimulation in bean (*Phaseolus vulgaris* L.) by *Trichoderma*. *Biol Control* 2009; 51: 409-16.
- [107] Vinale F, Sivasithamparam K, Ghisalberti EL, Ruocco M, Woo S, Lorito M. *Trichoderma* secondary metabolites that affect plant metabolism. *Nat Prod Commun* 2012; 7: 1545-50.
- [108] Cutler HG, Cox RH, Crumley FG, Cole PD. 6-Pentyl-a-pyrone from *Trichoderma harzianum*: its plant growth inhibitory and antimicrobial properties. *Agr Biol Chem Tokyo* 1986; 50: 2943-5.
- [109] Cutler HG, Himmetsbach DS, Arrendale RF, Cole PD, Cox RH. Koninginin A: a novel plant regulator from *Trichoderma koningii*. *Agr Biol Chem Tokyo* 1989; 53: 2605-11.
- [110] Parker SR, Cutler HG, Jacyno JM, Hill RA. Biological activity of 6-pentyl-2H-pyran-2-one and its analogs. *J Agri Food Chem* 1997; 45: 2774-6.
- [111] Parker SR, Cutler HG, Schreiner PR. Koninginin C: a biologically active natural product from *Trichoderma koningii*. *Biosci Biotech Bioch* 1995; 59: 1126-7.
- [112] Macías FA, Varela RM, Simonet AM, *et al.* Bioactive carotenes from *Trichoderma virens*. *J Nat Prod* 2000; 63: 1197-200.
- [113] Cutler HG, Jacyno JM. Biological activity of (-)-harzianopyridone isolated from *Trichoderma harzianum*. *Agr Biol Chem Tokyo* 1991; 55: 2629-31.
- [114] Cutler HG, Jacyno JM, Phillips RS, von Tersch RL, Cole PD, Montemurro N. Cyclonerodiol from a novel source, *Trichoderma koningii*: plant growth regulatory activity. *Agr Biol Chem Tokyo* 1991; 55: 243-4.
- [115] Marfori EC, Kajiyama S, Fukusaki E, Kobayashi A. Trichosetin, a novel tetramic acid antibiotic produced in dual culture of *Trichoderma harzianum* and *Catharanthus roseus* callus. *Z Naturforsch C* 2002; 57: 465-70.
- [116] Marfori EC, Kajiyama S, Fukusaki E, Kobayashi A. Phytotoxicity of the tetramic acid metabolite trichosetin. *Phytochemistry* 2003; 62: 715-21.
- [117] Vedonder RF, Tjarks LW, Rohwedder WK, Burmeister HR, Laugal JA. Equisetin, an antibiotic from *Fusarium equiseti* NRRL 5537, identified as a derivative of n-methyl-2,4-pyrrolidone. *J Antibiotics* 1979; 32: 759-61.

- [118] Kim H, Vinale F, Ghisalberti EL, *et al.* An antifungal and plant growth promoting metabolite from a sterile dark ectotrophic fungus. *Phytochemistry* 2006; 67: 2277-80.
- [119] Vinale F, Ghisalberti EL, Sivasithamparam K, *et al.* Factors affecting the production of *Trichoderma harzianum* secondary metabolites during the interaction with different plant pathogens. *Let Appl Microbiol* 2009; 48: 705-11.
- [120] Mukherjee PK, Horwitz BA, Kenerley CM. Secondary metabolism in *Trichoderma* - a genomic perspective. *Microbiology* 2012; 158: 35-45.

Received: November 21, 2013

Revised: January 07, 2014

Accepted: March 31, 2014

© Vinale *et al.*; Licensee *Bentham Open*.

This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>), which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.