

# Serum IL-6 Level in Breast Cancer Surgery: Evaluating the Addition of Hydrocortisone to Two Anesthetic Regimens

Sherif Abdelhamid<sup>1,\*</sup>, Ahmed Talha<sup>2</sup>, Salwa Hamdy<sup>3</sup>, Ashraf Arafat Abdelhalim<sup>4</sup> and Mohamed Elakany<sup>1</sup>

<sup>1</sup>Department of Anesthesia, Medical Research Institute, Alexandria University, Alexandria, Egypt

<sup>2</sup>Department of Surgery, Medical Research Institute, Alexandria University, Alexandria, Egypt

<sup>3</sup>Chemical Pathology, Medical Research Institute, Alexandria University, Alexandria, Egypt

<sup>4</sup>Department of Anesthesia, Alexandria University, Alexandria, Egypt

**Abstract:** *Background:* Interleukin (IL-6) and C-reactive protein (CRP) are found to be elevated in various inflammatory and malignant diseases and their levels are found to correlate with the extent of the disease. We evaluated the role of adding hydrocortisone to two different anesthetic regimens, on the level of IL-6, CRP and cortisol levels following breast cancer surgery.

*Patients and Methods:* In this prospective, randomized, controlled trial, 124 ASA I-II, female patients, aged 30-50 years, undergoing conservative breast cancer surgery were randomly assigned to one of four equal groups receiving either standard general anesthesia and two doses of hydrocortisone (Group GH, n=31), thoracic paravertebral block and two doses of hydrocortisone (Group PH, n=31), standard general anesthesia with no hydrocortisone (Group G, n=31), or thoracic paravertebral block with no hydrocortisone (Group P, n=31). IL-6 was measured at three time points: before operation, 6 and 12 hours postoperatively. CRP and cortisol were measured preoperatively and 6 hours postoperatively.

*Results:* On comparing group PH and GH, there was significant decrease in IL-6 level in group PH compared to group GH at 6 hour ( $122.1 \pm 21.2$  vs  $135.8 \pm 29.8$  pg/dl), but insignificant difference at 24 hours ( $107.9 \pm 21.6$  vs  $106.8 \pm 15.9$  pg/dl). CRP showed significant decrease in the postoperative reading in group PH compared to group GH ( $1.63 \pm 0.32$  vs  $1.91 \pm 0.43$  mg/l), and also group PH showed significant decrease compared to the control group P ( $1.63 \pm 0.32$  vs  $2.2 \pm 0.54$ ).

*Conclusion:* addition of hydrocortisone to general anesthesia or thoracic paravertebral block attenuated production of IL-6 and CRP levels significantly postoperatively compared to either anesthetic regimen alone, but not the serum cortisol level, highlighting its role in modifying the stress response to surgery. However, the effect was more pronounced when combined with thoracic paravertebral block.

**Keywords:** Anesthetic, breast, cancer, hydrocortisone, IL-6, regimens.

## INTRODUCTION

The cytokines are low-molecular-weight heterogenous proteins which include interleukins and interferons. The main cytokines released are interleukin-1 (IL-1), tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) and Interleukin (IL-6) [1]. They have a major role in the inflammatory response to malignancy, trauma and surgery, inducing local and systemic effects which attempt to limit injury and seem to be directly involved in the symptomatology of sepsis, multiorgan failure and septic shock [2]. IL-6 is a sensitive, early marker of tissue damage, and the main proinflammatory cytokine responsible for inducing acute phase response [1, 3]. It has a

major impact on outcome following surgery as its levels are increased in breast cancer patient. Excessive IL-6, both circulating and within the breast tumor microenvironment, is associated with poor clinical outcomes in breast cancer [4]. IL-6 stimulates secretion of C-reactive protein (CRP) from the liver, which is now considered to be an important biomarker for proinflammatory status in several diseases including cancer [5]. Both IL-6 and CRP are found to be elevated in various inflammatory and malignant diseases and their levels are found to correlate with the extent of the disease [6]. Serum IL-6 is released within minutes by blood leukocytes and constantly detected in the peripheral blood; therefore, the increase of IL-6 may reflect the magnitude of surgical trauma. By 24 to 36 hours postoperatively, the levels of IL-6 in the plasma decrease to preoperative values because of attenuated production [7-10]. It is believed that it stimulate auto and paracrine stimulation of tumor cell

\*Address correspondence to this author at the Department of Anesthesia, Medical Research Institute, Alexandria University, Alexandria, Egypt; E-mail: [dr\\_shereif@hotmail.com](mailto:dr_shereif@hotmail.com)

proliferation, in addition of being an independent prognostic marker of survival in metastatic breast cancer patients [11]. In literature, there are few researches focusing on the impact of hydrocortisone on cytokine levels in humans. In a prior study, the use of hydrocortisone led to significant decrease of the inflammatory response among patients undergoing bilateral total knee replacement indicated by measuring IL-6 production [12]. This prospective, randomized, controlled study was performed to evaluate possible differences in the plasma inflammatory cytokine interleukin 6 (IL-6), the release of the stress hormones (cortisol), and C-reactive protein (CRP) by adding hydrocortisone to two different anesthetic regimens in a population of healthy female patients with cancer breast undergoing unilateral conservative breast cancer surgery.

## PATIENTS AND METHODS

After obtaining Institutional Review Board approval (Trial number: PACTR201304000509378) and written informed consent, this prospective, randomized, controlled study was conducted at Medical Research Institute Hospital, Alexandria, Egypt in the period from April 2013 to February 2014. The study enrolled 124 female patients, ASA physical status I or II aged 30 to 50 years, with cancer breast scheduled for unilateral conservative breast cancer surgery.

Patients with endocrine or metabolic disorders, diabetes, morbid obesity (Body Mass Index  $>40 \text{ kg/m}^2$  or  $35 \text{ kg/m}^2$  with co-morbidity), history of CNS diseases, known allergy to bupivacaine or any contraindication to paravertebral block, pregnancy, lactation, taking steroid medications and patients under current or recent medication affecting the sympathetic response or the hormonal secretion were excluded from the study.

All patients were subjected to a standard pre-operative evaluation. The selected patients were randomly allocated to one of the four equal groups by using sealed, opaque envelopes to receive standard general anesthesia and two doses of hydrocortisone (Group GH,  $n=31$ ), thoracic paravertebral block and two doses of hydrocortisone (Group PH,  $n=31$ ), standard general anesthesia with no hydrocortisone (Group G,  $n=31$ ), or thoracic paravertebral block with no hydrocortisone (Group P,  $n=31$ ).

In group (GH), on arrival at the operating theatre, routine standard monitoring was used for all patients, after preoxygenation using 100% Oxygen for 3 minutes, all patients received standard anesthetic techniques; anesthesia was induced by fentanyl  $1.5 \mu\text{g/kg}$ , and propofol  $2 \text{ mg/kg}$  followed by the administration of cisatracurium  $0.15 \text{ mg/kg}$  to facilitate orotracheal intubation. Anesthesia was maintained with isoflurane 1–1.5% in a mixture of air in oxygen (60%:40%), and patients received intermittent doses of fentanyl and cisatracurium. All patients received hydrocortisone  $100 \text{ mg}$  intravenously at induction and the same dose was repeated 2 hours later.

Patients in Group (PH) were sedated during performance of thoracic paravertebral block (TPVB) with propofol infusion in a dose of  $100 \text{ mcg/kg/min}$  using a syringe pump (Life Care5000 Infusion Set, Abbott, Ireland). Thoracic paravertebral blocks were then performed as described by

Katz [13]. Paravertebral injections were performed with the patient in lateral position (surgical side facing upward). Spinous processes of T3 to T6 were identified and a mark was taken corresponding to 2.5 cm lateral to them. After aseptic preparation of the skin, these points were infiltrated with lidocaine 1% 1–2 ml. An 18-gauge Epidural Tuohy needle was inserted perpendicular to the skin at T4 at the corresponding mark, when the tip of the needle touched the corresponding transverse process, the needle was withdrawn and redirected cephalad to slide over the transverse process to reach a depth of 4 cm as a maximum, piercing the superior costotransverse ligament and advanced into the paravertebral space. A loss of resistance to saline was usually encountered to confirm the space at this depth. After negative aspiration of blood, air or cerebrospinal fluid, 30 ml of bupivacaine 0.5% was injected; adequate block was confirmed by loss of temperature sensation over appropriate dermatomes, and by testing pinprick sensation in the surgical field before incision. All patients were contacted verbally and routine standard monitoring was used for all patients. After assuring adequate anesthesia and the patient was vitally stable, without any manifestations of local anesthetic toxicity, the surgery started. All patients received hydrocortisone  $100 \text{ mg}$  at start of block and the same dose was repeated two hours later. Two senior consultants performed the TPVBs to minimize variation in the technique.

Patients in group (G) received the same standard anesthetic techniques as group (GH) without administration of Hydrocortisone and this group was served as control group.

Patients in group (P) received the same thoracic paravertebral block technique as group (P) without administration of Hydrocortisone and this group was served as control group.

Postoperative pain scores in the first 24 hours were always below or equal 3 on visual analogue scale (VAS) in all studied groups. Infusion of 0.125 % bupivacaine in the paravertebral groups, while using 30 mg ketorolac/ 8 hours IV. in the general anesthesia groups and nalbuphine of 2 mg IV. As a rescue analgesic, to maintain VAS of  $\leq 3$ .

Venous blood samples were collected peripherally, from an antecubital vein of the arm contralateral to the surgery, using a small cannula. For each patient in all groups, peripheral blood samples for measuring IL-6 were collected at three time points: before operation, six hour after the end of the procedure, and twenty four hours after the procedure. IL-6 was measured in serum samples by enzyme linked immunosorbent assay using AviBion Human IL-6 ELISA KIT, FINLAND, The assay employed an antibody specific for human IL-6 coated on a 96-well plate. IL-6 in samples is captured by the antibody immobilized to the wells and by a biotinylated IL-6 specific detection antibody. After washing away unbound biotinylated antibody, HRP-conjugated streptavidin and TMB substrate solution were added to the wells resulting in color development proportional to the amount of IL-6 bound. The intensity of the color was measured at 450 nm. The mean of reagent blank absorbance values were calculated and subtracted from all test well (standard and samples). A calculated result against standard curve was drawn [14]. Serum and plasma samples were



9.54±1.6 vs 20.93±1.74, 9.50±1.5 vs 20.72±1.60 in groups GH, PH, G and P respectively ( $P < 0.05$ , Table 3), however, the difference between groups was statistically insignificant ( $P > 0.05$ , Table 3).

All four groups showed significant increase in CRP postoperatively compared to baseline value ( $P < 0.05$ ,

Table 4), however, the CRP value was significantly less in group PH compared to group GH postoperatively (1.63±0.32 vs 1.91±0.43mg/l, respectively,  $P < 0.05$ , Table 4). The CRP values was significantly less in both group PH and group GH compared to their controls; (1.91±0.43 vs 2.23±0.40 mg/l) and (1.63±0.32 vs 2.2±0.54) respectively,  $P < 0.05$ , Table 4).

**Table 1. Patients characteristics.**

Personal characteristics	Group (GH) (n=31)	Group (PH) (n=31)	Control (G) (n=31)	Control (P) (n=31)
Age (years)				
Min-Max	30-49	33-48	31-45	34-50
Mean±SD	41.1±3.2	42.6±2.9	39.8±2.8	44.5±5.1
BMI (Kg/m <sup>2</sup> )				
Min-Max	26.6-34.4	28.1-35.8	27.4-34.9	26.2-35.4
Mean±SD	31.2±4.7	32.6±3.9	31.8±2.7	32.1±4.5
Duration of surgery (min)				
Min-Max	70-90	70-85	65-90	70-95
Mean±SD	81.1±3.1	79.7±3.3	79.5±5.9	82.0±6.4
ASA				
ASA I	11 (35.5%)	10 (32.3%)	12 (38.7%)	7 (22.6%)
ASA II	20 (64.5%)	21 (67.7%)	19 (61.3%)	24 (77.4%)

ASA = American Society of Anesthesiologists. Values are presented as means (SD), or number of patients (n) and percentages (%).

**Table 2. Interleukin- 6 measured pre and postoperatively among the studied groups.**

Interleukin 6 (pg/dl)	Group (GH) (n=31)	Group (PH) (n=31)	Control (G) (n=31)	Control (P) (n=31)	Sig 1	Sig 2	Sig 3
Preoperative							
Min-Max	35-130	45-130	53-120	50-120	t=0.711	t=0.393	t=0.42
Mean±SD	86.3±20.9	89.8±18.4	88.0±13.9	87.7±13.9	P=0.48	P=0.696	P=0.676
6 hours post-operative							
Min-Max	80-200	70-160	100-280	100-280	t=2.134	t=2.453	t=4.461
Mean±SD	135.8±29.8	122.1±21.2	158.1±40.8	156.9±39.6	P=0.037*	P=0.017*	P<0.0001*
24 hours post-operative							
Min-Max	55-150	80-140	95-260	95-260	t=0.172	t=4.912	t=5.46
Mean±SD	107.6±21.9	106.8±15.9	147.7±39.8	146.9±39.3	P=0.864	P<0.0001*	P<0.0001*
<i>Sig (Pre/6 hrs Postoperative)</i>	[P<0.0001*]	[P<0.0001*]	[P<0.0001*]	[P<0.0001*]			
<i>Sig (Pre/24 hrs Postoperative)</i>	[P<0.0001*]	[P<0.0001*]	[P<0.0001*]	[P<0.0001*]			
<i>Sig (Post 6 hrs/24 hrs Postoperative)</i>	[P<0.0001*]	[P<0.0001*]	[P<0.0001*]	[P<0.0001*]			

Sig1: Significance between Group (GH) and Group (PH)

Sig2: Significance between Group (GH) and Control (G) Group

Sig3: Significance between Group (PH) and Control (P) Group

t: t-test [P]: P for Paired t-test \*significant at  $P \leq 0.05$

Fig. (2). Interleukin- 6 measured pre and postoperatively among the studied groups.

Table 3. Serum cortisol level measured pre and postoperatively among the studied groups.

Serum cortisol level	Group (GH) (n=31)	Group (PH) (n=33)	Control (G) (n=31)	Control (P) (n=31)
Preoperative				
Min-Max	9.1-10.0	9.3-10.1	8.9-10.2	8.8-10.0
Mean±SD	9.41±1.40	9.52±1.41	9.54±1.6	9.50±1.5
Post-operative				
Min-Max	18.8-22.0	18.5-23.4	19.0-22.4	18.1-22.3
Mean±SD	20.11±1.72*	20.60±1.61*	20.93±1.74*	20.72±1.60*

\* P<0.05 significant compared to baseline.

Table 4. CRP level measured pre and postoperatively among the studied groups.

CRP (mg/L)	Group (GH) (n=31)	Group (PH) (n=33)	Control (G) (n=31)	Control (P) (n=31)
Preoperative				
Min-Max	0.7-1.8	0.8-1.9	0.9-1.8	0.6-1.9
Mean±SD	1.24±0.31	1.32±0.37	1.21±0.30	1.23±0.44
Post-operative				
Min-Max	1.2-3.0	1.2-2.6	1.8-3.2	1.8-3.4
Mean±SD	1.91±0.43*‡	1.63±0.32*†	2.23±0.40*	2.2±0.54*

\* P<0.05 significant compared to baseline, † P=0.001 significant compared to group GH,

‡ P<0.0001 significant compared to group G, P<0.0001 significant compared to group P.

## DISCUSSION

This prospective, randomized, controlled study revealed that the addition of two doses of hydrocortisone to either general anesthesia or thoracic paravertebral block attenuated production of IL-6 significantly at 6 and 24 hours postoperatively compared to control groups, demonstrating the role of hydrocortisone in reducing the IL-6 level. In animal models, Lodge *et al.* [16] showed that the use of glucocorticoids can reduce the interleukins 6 and 8, tumor necrosis factor-(TNF- $\alpha$ ) and increase anti-inflammatory cytokines such as IL-10, IL-4.

Elysee *et al.* [12] reported that, the use of hydrocortisone (100 mg) at 2 doses 8 hrs apart significantly decreased the inflammatory response in patients undergoing bilateral total knee replacement as the levels of IL-6 were 40% lower in the study group compared to placebo group. Moreover, the range of motion gained on discharge was also greatest in the study group. In another study, it has been suggested that combined preoperative and intraoperative steroid administration attenuates the inflammatory response more effectively than intraoperative steroid administration alone [17].

In our study, the baseline level of IL-6 was elevated in all groups. Hussien MZ *et al.* [18] in their study showed that the baseline IL-6 level is highly increased in breast cancer patients, and that serum levels of IL-6 (mean =111.38 pg/ml) was significantly increased in breast cancer patients as compared to controls (mean 1.75 pg/ml). Additionally, Dethlefsen *et al.* [19] illustrated in their study that high circulating IL-6 levels are correlated with a poor prognosis in breast cancer patients. Surgical stress is known to induce an increase in serum levels of IL-6. However, different regimens of anesthesia are associated with differences in cytokine secretion pattern.

A combination of hydrocortisone and general anesthesia provided significantly less attenuation of production of IL-6 compared to combination of hydrocortisone with thoracic paravertebral block at 6 hours but not 24 hours postoperatively. These results indicate that TPVB seems to better attenuate the production of IL-6 compared to general anesthesia. However, in the current study, TPVB was not continued postoperatively by using a catheter technique in the paravertebral space. This may explain insignificant differences between groups 24-h post-operatively. Many anesthetic techniques are used to decrease immune responses to surgeries, and it has been reported that the stress response to surgery, is attenuated by regional anesthesia, including paravertebral anesthesia by blocking pain pathways [20-22].

The results of the current study supported the findings reached by Cuellar *et al.* [23] who measured the proinflammatory cytokines in subjects undergoing laparoscopic cholecystectomy, receiving either regional or general anesthesia, comparing the 2 groups, they argued that proinflammatory cytokines was significantly higher in the general anesthesia group. In another study by, Hadimioglu *et al.* [24] they showed that IL-6 levels increased significantly at all the measured times compared with the baselines level in both general anesthesia group, and combined epidural

with general anesthesia group. On comparing the 2 groups, they observed that the increase was greater in the general anesthesia compared with the combined group at all measured times.

Experimental studies in rats showed that surgical stress is attenuated better by the addition of spinal blockade to general halothane anesthesia than by general anesthesia alone and that, consequently, natural killer cell function is better preserved and markedly attenuates the promotion of metastasis by surgery [25]. A retrospective analysis of 129 patients undergoing mastectomy and axillary lymph node dissection for breast cancer suggested a lower risk for cancer recurrence or metastasis during the initial years of follow-up in patients having surgery with paravertebral block combined with a general anesthesia compared with patients with general anesthesia alone [26].

Our study showed that the postoperative increase of C-reactive protein concentrations was significantly less with combination of hydrocortisone with thoracic paravertebral block compared to combination of hydrocortisone and general anesthesia. Moreover, addition of hydrocortisone to either group provided significantly less increase compared to control groups highlighting that addition of hydrocortisone to anesthetic regimens could possibly decrease stress response markers which was more pronounced when added to thoracic paravertebral block. This result confirms previous clinical data, O'Riain *et al.* [21] showed that CRP as a marker of stress response were reduced in general anesthesia-paravertebral block group for breast cancer surgery compared to general anesthesia alone group (2.6 $\pm$ 1.5 vs 5.7 $\pm$ 2.9 nmol/L). In another study, Rodriguez *et al.* [27] showed that thoracic epidural anesthesia significantly attenuated the CRP level.

Although, the postoperative serum cortisol level was significantly increased in all groups compared to baseline value, the difference between groups was insignificant. Conflicting results have been reported by Wedad *et al.* [28] who found that the serum cortisol level postoperatively was significantly lower in paravertebral group. In other study, Ibrahim *et al.* [29] found that the paravertebral group had significant lower values of postoperative serum cortisol compared to thoracic epidural group. In other report, Aggo *et al.* [30] compared plasma cortisol level in epidural versus general anesthesia groups, and found significant decrease in epidural group in all measured times. Our study is not quite similar with previous studies because of the variations in study design; the dosage, timing of hydrocortisone administration and timing of serum cortisol level measurement.

Limitations of the study include that; first, we measured only one inflammatory cytokine, future studies would benefit from assessments of more cytokines. Second, serum cortisol level would need more monitoring to trace the changes after I.V hydrocortisone injection. Third, we did not measure any clinical outcomes; therefore further clinical trials would be required to ascertain whether the differences in cytokine levels affect tumor recurrence, wound healing and patient survival. Finally, more studies on larger scale are warranted with possibly more doses of hydrocortisone.

## CONCLUSION

We found that of addition of hydrocortisone to either general anesthesia or thoracic paravertebral block attenuated production of IL-6 and CRP levels significantly postoperatively compared to either anesthetic regimen alone, but not the serum cortisol level, highlighting its role in modifying the stress response to surgery. However, the effect was more pronounced when combined with thoracic paravertebral block, it is possible that TPVB may offer advantages in terms of inflammatory effects compared to general anesthesia.

## CONFLICT OF INTEREST

The authors confirm that this article content has no conflict of interest.

## ACKNOWLEDGEMENTS

The authors are grateful to the staff members of operating room, Medical Research Institute Hospital, Alexandria University, Alexandria, Egypt.

## REFERENCES

- [1] Desborough JP. The stress response to trauma and surgery. *Br J Anaesth* 2000; 85: 109-17.
- [2] Sheeran P, Hall GM. Cytokines in anaesthesia. *Br J Anaesth* 1997; 78: 201-19.
- [3] Helmy SA, Wahby MA, El-Nawaway M. The effect of anaesthesia and surgery on plasma cytokine production. *Anaesthesia* 1999; 54(8): 733-8.
- [4] Bachelot T, Ray-Coquard I, Menetrier-Caux C, *et al.* Prognostic value of serum levels of interleukin 6 and of serum and plasma levels of vascular endothelial growth factor in hormone-refractory metastatic breast cancer patients. *Br J Cancer* 2003; 88: 1721-6.
- [5] Erlinger TP, Platz EA, Rifai N, *et al.* C-reactive protein and the risk of incident colorectal cancer. *JAMA* 2004; 291: 585-90.
- [6] Ravishankaran P, Karunanithi R. Clinical significance of preoperative serum interleukin-6 and C-reactive protein level in breast cancer patients. *World J Surg Oncol* 2011; 9: 18.
- [7] Holzheimer RG, Steinmetz W. Local and systemic concentrations of proand anti-inflammatory cytokines in human wounds. *Eur J Med Res* 2000; 5(8): 347-55.
- [8] Notarnicola A, Moretti L, Tafuri S, *et al.* Postoperative pain monitor after total knee replacement. *Musculoskelet Surg* 2011; 95(1): 19-24.
- [9] Pandazi A, Kapota E, Matsota P, *et al.* Preincisional versus postincisional administration of parecoxib in colorectal surgery: effect on postoperative pain control and cytokine response. A randomized clinical trial. *World J Surg* 2010; 34(10): 2463-9.
- [10] Lin E, Calvano SE, Lowry SF. Inflammatory cytokines and cell response in surgery. *Surgery* 2000; 127(2): 117-26.
- [11] Salgado R, Junius S, Benoy I, *et al.* Circulating IL-6 predicts survival in patients with metastatic breast cancer. *Int J Cancer* 2003; 103: 642-6.
- [12] Elysee J, Kethy M, Lipnitsky, *et al.* Use of low dose steroids in decreasing cytokine release during bilateral total knee replacement. *Reg Anesth Pain Med* 2011; 36: 36-40.
- [13] Katz J. Atlas of regional anesthesia. 2<sup>nd</sup> ed. Norwalk: Appleton & Lange 1994.
- [14] Allen MJ1, Myer BJ, Khokher AM, *et al.* Pro-inflammatory cytokines and the pathogenesis of Gaucher's disease: increased release of interleukin-6 and interleukin-10. *QJM* 1997; 90(1): 19-25.
- [15] Prieto B, Miguel D, Costa M, *et al.* New quantitative electrochemiluminescence method (ECLIA) for interleukin-6 (IL-6) measurement. *Clin Chem Lab Med* 2010; 48(6): 835-8.
- [16] Lodge AJ, Chai PJ, Daggett CW, *et al.* Methylprednisolone reduces the inflammatory response to cardiopulmonary bypass in neonatal piglets: Timing of dose is important. *J Thoracic Cardiovasc Surg* 1999; 117(3): 515-22.
- [17] Schroeder VA, Pearl JM, Schwartz SM, *et al.* Combined steroid treatment for congenital heart surgery improves oxygen delivery and reduces postbypass inflammatory mediator expression. *Circulation* 2003; 107(22): 2823-8.
- [18] Hussein MZ, Al Fikky A, Abdel Bar I, *et al.* Serum IL-6 and IL-12 levels in breast cancer patients. *Egypt J Immunol* 2004; 11(2): 165-70.
- [19] Dethlefsen C, Højfeldt G, Hojman P. High circulating IL-6 levels are correlated with a poor prognosis in breast cancer patients. *Breast Cancer Res Treat* 2013; 138(3): 657-64.
- [20] Buggy DJ, Smith G. Epidural anaesthesia and analgesia: better outcome after major surgery? Growing evidence suggests so. *BMJ* 1999; 319: 530-1.
- [21] O'Riain SC, Buggy DJ, Kerin MJ, *et al.* Inhibition of the stress response to breast cancer surgery by regional anesthesia and analgesia does not affect vascular endothelial growth factor and prostaglandin E2. *Anesth Analg* 2005; 100: 244-9.
- [22] Sultan SS. Paravertebral block can attenuate cytokine response when it replaces general anesthesia for cancer breast surgeries. *Saudi J Anaesth* 2013; 7(4): 373-7.
- [23] Cuellar A, Ramos J, Flores G, *et al.* Comparison of two anesthetic techniques on the plasma level of inflammatory markers. *Cir Ciruj* 2007; 75: 99-105.
- [24] Hadimioglu N, Ulugol H, Akbas H, *et al.* Combination of epidural anesthesia and general anesthesia attenuates stress response to renal transplantation surgery. *Transplant Proc* 2012; 44(10): 2949-54.
- [25] Bar-Yosef S, Melamed R, Page GG, *et al.* Attenuation of the tumor-promoting effect of surgery by spinal blockade in rats. *Anesthesiology* 2001; 94: 1066-73.
- [26] Exadakylos AK, Buggy DJ, Moriarty DC, *et al.* Can anesthetic technique for primary breast cancer surgery affect recurrence or metastasis? *Anesthesiology* 2006; 105: 660-4.
- [27] Rodriguez MA, Gonzalo IA, Alvarez EV, *et al.* Thoracic epidural anesthesia attenuates CRP levels in patients undergoing elective coronary artery bypass graft surgery with cardiopulmonary bypass. *Minerva Anestesiol* 2008; 74: 6619-26.
- [28] Wedad M, Zaki K, Haleem M. The effect of addition of wound infiltration with local anaesthetic to interpleural block on post-thoracotomy pain, pulmonary function and stress response in comparison to epidural and paravertebral block. *Eg J Anaesth* 2004; 20: 67-72.
- [29] Ibrahim A, Mamdouh N. Comparison between continuous thoracic epidural block and continuous thoracic paravertebral block for thoracotomy pain relief. *Ain Shams J Anesth* 2009; 2: 16-26.
- [30] Aggo AT, Fyneface-Ogan S, Mato CN. The differential impact of two anesthetic techniques on cortisol levels in Nigerian surgical patients. *Nigerian J Clin Pract* 2012; 15: 68-74.

Received: February 03, 2015

Revised: March 01, 2015

Accepted: March 17, 2015

© Abdelhamid *et al.*; Licensee Bentham Open.

This is an open access article licensed under the terms of the (<https://creativecommons.org/licenses/by/4.0/legalcode>), which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.